

COMBINATION SHEEP DRENCH



WARNING

Dangerous to the Environment.

Very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment.

PLEASE READ LABEL COMPLETELY BEFORE USE

PRECAUTIONS:

Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose or dose more frequently than three weekly. In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.

HANDLING PRECAUTIONS:

May be harmful if swallowed. Repeated exposure may cause skin allergy and mild skin irritation. Albendazole and levamisole hydrochloride may possibly cause damage to genetic material and may affect development and/or reproduction. Levamisole hydrochloride may possibly affect blood and the haemopoietic system. Avoid skin contact. Wear gloves when decanting. Do not eat, drink or smoke when using this product. Avoid release to the environment.

FIRST AID:

If swallowed, Do NOT induce vomiting. For advice contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately. If skin or hair contact occurs, remove contaminated clothing and wash skin thoroughly. If splashed in eyes, flush immediately with copious amounts of water.

EMERGENCY RESPONSE

In case of spillage wear appropriate clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

INDICATIONS

Contains albendazole (23.8 mg/mL), levamisole HCl 37.5 mg/mL, selenium 0.5 mg/mL, cobalt 0.3 mg/mL, copper 2.1 mg/mL, zinc 0.6 mg/mL. For the treatment of tapeworms and liver fluke and the following susceptible adult and immature stages of gastrointestinal roundworms and lungworms including benzimidazole and levamisole resistant strains: *Haemonchus contortus*, *Ostertagia* spp., *Trichostrongylus* spp., *Nematodirus* spp., *Cooperia* spp., *Strongyloides papillosus*, *Bunostomum trigonocephalum*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Trichuris ovis* (adults only), *Moniezia expansa* (segments only), *Dictyocaulus* spp., *Muellerius capillaris*, *Liver fluke* (adults only). Also reduces the output of roundworm and fluke.

RESISTANCE MANAGEMENT

Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinary practitioner for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

DIRECTIONS FOR USE

By law, the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress in the treated animal when using the product in a manner other than as recommended on the label.

Shake well before use. Combination Sheep Drench is a ready to use anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment weigh a representative sample of sheep to determine the weight of the heaviest animal. Where there is a large variation in liveweights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band. Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

Dose Chart

Dose rate sheep: 1 mL/5 kg (equivalent to 4.75 mg/kg albendazole and 7.5 mg/kg levamisole HCl.)

ANIMAL WEIGHT	DOSE	NO. OF DOSES PER 5L PACK
15 kg	3 mL	1666
16 - 20 kg	4 mL	1250
21 - 25 kg	5 mL	1000
26 - 30 kg	6 mL	833
31 - 35 kg	7 mL	714
36 - 40 kg	8 mL	625
41 - 45 kg	9 mL	555
46 - 50 kg	10 mL	500
51 - 55 kg	11 mL	454
56 - 60 kg	12 mL	416
61 - 65 kg	13 mL	384
66 - 70 kg	14 mL	357

Over 70 kg dose at the rate of 1 mL/5 kg.

It is an offence under the law to sell produce from animals treated with this product with residues that exceed the maximum residue levels in the New Zealand (Maximum Residue of Agriculture Compounds) Food Standard.

WITHHOLDING PERIODS

Meat: Sheep producing meat or offal for sale for human consumption must not be sold for slaughter either during treatment or within **10 days** of the last treatment.

Milk: Milk intended for sale for human consumption must be discarded during treatment and for not less than **35 days** following the last treatment.

STORAGE

Store locked up in original container, tightly closed, in cool, dry conditions. Protect from light. Avoid temperatures below 0° and above 30°C.

Registered pursuant to the ACVM Act 1997, No. A9764.
See www.foodsafety.govt.nz for registration conditions.



Registered to:
Seneca Holdings Limited
0800 11 8888

Batch No.:

Expiry date:

